



2009 WORLD VOICE DAY

WORLD VOICE DAY

APRIL 16th

Laryngeal (Voice Box) Cancer

Laryngeal cancer is not as well known by the general public as some other types of cancer, yet it is not a rare disease. The American Cancer Society estimates that in the United States there were 12,250 new cases of laryngeal cancer diagnosed in 2008, and 3,670 deaths from the disease. Even for survivors, the consequences of laryngeal cancer can be severe with respect to voice, breathing, or swallowing. It is fundamentally a preventable disease though, since the primary risk factors for laryngeal cancer are associated with behaviors that can be modified.

Risk factors associated with laryngeal cancer

Development of laryngeal cancer is a process that involves many factors, but approximately 90 percent of head and neck cancers occur after exposure to known carcinogens (cancer-causing substances). Chief among these factors is tobacco. Over 90 percent of laryngeal cancers are a type of cancer called squamous cell carcinoma (SCCA), and over 95 percent of patients with laryngeal SCCA are smokers. Smoking contributes to cancer development by causing mutations or changes in genes, impairing clearance of carcinogens from the respiratory tract, and decreasing the body's immune response.

- **Tobacco** use is measured in pack-years, where one pack per day for one year is considered one pack-year. Two pack-years is defined as either one pack per day for two years, or two packs per day for one year (Longer terms of pack years are determined using a similar ratio). Depending upon the number of pack-years smoked, studies have reported that smokers are about 5 to 35 times more likely to develop laryngeal cancer than non-smokers. It does seem that the duration of tobacco exposure is probably more important overall as a cancer-causing effect, than the intensity of the exposure.

"Listen to your voice, it may be telling you something!"



AMERICAN ACADEMY OF
OTOLARYNGOLOGY-
HEAD AND NECK SURGERY

Empowering otolaryngologist—head and neck surgeons to deliver the best patient care

1650 Diagonal Road, Alexandria, Virginia 22314-2857 U.S.A. 1-703-836-4444 1-703-683-5100 fax www.entnet.org

- **Alcohol** is another important risk factor for laryngeal cancer, and acts as a promoter of the cancer causing process. The major clinical significance of alcohol is that it increases the effects of tobacco. That is, people who smoke and drink alcohol have a combined risk that is greater than the sum of the individual risks. The American Cancer Society recommends that those who drink alcoholic beverages should limit the amount of alcohol they consume, with one drink per day considered a limited alcohol exposure.
- Other risk factors for laryngeal cancer include certain viruses, such as the human papilloma virus (HPV), and possibly, acid reflux. Vitamin A and beta-carotene may play a protective role.

Signs and symptoms of laryngeal cancer

These include:

- Progressive or persistent hoarseness
- Difficulty swallowing
- Persistent sore throat or pain with swallowing
- Difficulty breathing
- Pain in the ear
- A lump in the neck

Anyone with these signs or symptoms should be evaluated by an otolaryngologist (ear, nose, and throat doctor). This is particularly important for people with risk factors for laryngeal cancer.

Treatment of laryngeal cancer

The primary treatment options for laryngeal cancer include surgery, radiation therapy, chemotherapy, or a combination of these treatments. Remember that this is a preventable disease in the vast majority of cases, because the main risk factors are associated with modifiable behaviors. Do not smoke and do not abuse alcohol.

For more information on vocal health, visit www.entnet.org/worldvoiceday.