



## **Clinical Indicators: Parotidectomy**

<b><u>Procedure</u></b>	<b>CPT</b>	<b>Days<sup>1</sup></b>
Excision of parotid tumor or parotid gland; lateral lobe, without nerve dissection	42410	90
Excision of parotid tumor or parotid gland; lateral lobe, with dissection and preservation of facial nerve	42415	90
Excision of parotid tumor or parotid gland; total, with dissection and preservation of facial nerve	42420	90
Excision of parotid tumor or parotid gland; total; en bloc removal with sacrifice of facial nerve	42425	90
Excision of parotid tumor or parotid gland; total, with unilateral radical neck dissection	42426	90

<b><u>Related Procedures</u></b>	<b>CPT</b>	<b>Days<sup>1</sup></b>
Drainage of abscess; parotid, simple	42300	10
Drainage of abscess, parotid, complicated	42305	90
Sialolithotomy; parotid, uncomplicated, intraoral	42330	10
Sialolithotomy; parotid, extraoral or complicated intraoral	42340	90
Biopsy of salivary gland, needle	42400	0
Biopsy of salivary gland, incisional	42405	10
Unlisted procedure, salivary glands or ducts	42699	

## **Indications**

### **1. History (one or more required)**

- a) Parotid mass.
- b) History of radiation to the neck.
- c) Chronic parotitis.
- d) A neck mass with histologic findings of metastatic parotid tumor.
- e) Parotid duct stone.
- f) Malignancy of overlying skin extending into parotid
- g) Malignancy metastatic to parotid.

---

<sup>1</sup> RBRVS Global Days



## **2. Related Symptoms**

- a) Facial nerve paralysis.
- b) Pain of parotid region.

## **3. Physical Examination** (required)

- a) Complete physical examination of the head and neck with emphasis on inspection and palpation of the parotid gland, oropharynx and neck.
- b) Examination of facial nerve function.

## **4. Tests** (required)

- a) Pre-operative tests as required by institutional guidelines.

## **5. Tests** (optional)

- a) Fine needle aspiration biopsy.
- b) Ultrasonography.
- c) CT scan of neck.
- d) MRI of neck.
- e) Sialogram
- f) Technetium-99 radionuclide imaging

## **Postoperative Observations**

- a) Facial nerve function
- b) Bleeding - check for expanding hematoma; notify surgeon

## **Outcome Review**

### **1. One Week**

- a) Facial nerve function - present in all branches of facial nerve?  
Need for protection of eye from drying?
- b) Wound healing well - infection or fistula?
- c) Pathology report - compare with pre-operative needle aspiration report if any. Are margins clear? Benign or malignant?
- d) Discuss with patient any ear numbness and/or gustatory sweating of facial skin.

### **2. Beyond One Month**

- a) If malignant - what type? Need for additional imaging? Is additional therapy indicated?
- b) If benign with tumor at margin - need for additional therapy?



**AMERICAN ACADEMY OF  
OTOLARYNGOLOGY–  
HEAD AND NECK SURGERY**

- c) Facial nerve function - present in all branches of nerve? Need for protection of eyes from drying? Need for additional rehabilitation?
- d) Gustatory sweating - need for treatment?
- e) e) Incision healed? Fistula present - management needed?

**3. Beyond One Year**

- a) Malignancy or incompletely excised benign tumor - need for imaging?
- b) Facial nerve function - need for rehabilitation?

**Associated ICD-9 Diagnostic Codes** (Representative, but not all-inclusive codes)

- 072.9 Mumps without mention of complication (Epidemic parotitis, infectious parotitis)
- 142.1-  
142.8 Malignant neoplasm of major salivary glands
- 142.0 Parotid gland
- 210.2 Benign neoplasm of major salivary glands
- 228.00-  
228.09 Hemangioma, any site
- 228.1 Lymphangioma, any site
- 230.0 Carcinoma in situ of lip oral cavity and pharynx
- 235.1 Neoplasm of uncertain behavior of digestive and respiratory systems
- 235.0 Major salivary glands
- 351.0 Bell's palsy
- 527.1 Hypertrophy
- 527.2 Sialoadenitis
- 527.5 Sialolithiasis
- 527.6 Mucocele
- 527.7 Disturbance of salivary secretion
- 527.8 Other specified diseases of the salivary glands
- 527.9 Unspecified disease of the salivary glands

**Associated ICD-10-CM Diagnostic Codes** (Representative, but not all-inclusive codes)

- B26.9 Mumps without complication
- C07 Malignant neoplasm of parotid gland
- D11.9 Benign neoplasm of major salivary gland, unspecified
- D11.0 Benign neoplasm of parotid gland



**AMERICAN ACADEMY OF  
OTOLARYNGOLOGY–  
HEAD AND NECK SURGERY**

- D11.7 Benign neoplasm of other major salivary glands
- D18.00 Hemangioma unspecified site
- D18.1 Lymphangioma, any site
- D00.00 Carcinoma in situ of oral cavity, unspecified site
- D00.01 Carcinoma in situ of labial mucosa and vermilion border
- D00.02 Carcinoma in situ of buccal mucosa
- D00.03 Carcinoma in situ of gingiva and edentulous alveolar ridge
- D00.04 Carcinoma in situ of soft palate
- D00.05 Carcinoma in situ of hard palate
- D00.06 Carcinoma in situ of floor of mouth
- D00.07 Carcinoma in situ of tongue
- D00.08 Carcinoma in situ of pharynx
- D37.030 Neoplasm of uncertain behavior of the parotid salivary glands
- D37.031 Neoplasm of uncertain behavior of the sublingual salivary glands
- D37.032 Neoplasm of uncertain behavior of the submandibular salivary glands
- D37.039 Neoplasm of uncertain behavior of the major salivary glands, unspecified
- G51.0 Bell's palsy
- K11.0 Atrophy of salivary gland
- K11.1 Hypertrophy of salivary gland
- K11.20 Sialoadenitis, unspecified
- K11.21 Acute sialoadenitis
- K11.22 Acute recurrent sialoadenitis
- K11.23 Chronic sialoadenitis
- K11.3 Abscess of salivary gland
- K11.4 Fistula of salivary gland
- K11.5 Sialolithiasis
- K11.6 Mucocele of salivary gland
- K11.7 Disturbances of salivary secretion
- R68.2 Dry mouth, unspecified
- K11.8 Other diseases of salivary glands
- K11.9 Disease of salivary gland, unspecified

**Additional Information**

Assistant Surgeon -- Varies

Supply Charges -- N

Prior Approval - N/A

Anesthesia Code(s) -- 00100; 00300

**Patient Information**

Parotidectomy is a surgical operation to remove a large salivary gland (the parotid gland) located in front and just below the ear. The most common reasons for removal of all or part of this gland are a mass in the gland, chronic infection of the gland, or obstruction of the saliva outflow from the gland causing chronic enlargement of the gland. Masses in the parotid are most commonly



**AMERICAN ACADEMY OF  
OTOLARYNGOLOGY–  
HEAD AND NECK SURGERY**

benign, but about 20% are malignant. The physician will discuss with you the need for parotidectomy based on your medical history, the results of a physical examination of the head and neck, and results of other tests if indicated. The most common tests to determine whether a parotidectomy is necessary include a fine needle aspiration biopsy (withdrawing a small amount of fluid from the parotid to see if malignant cells are present), CT scan (an x-ray test that helps to determine the size and position of the parotid tissues), and MRI (an imaging test that does not use x-rays and helps to determine the size and position of parotid tissues). In some cases no additional testing may be needed prior to surgery.

The procedure is usually done under general anesthesia. The amount of parotid gland to be removed is often determined at the time of surgery based on the size and location of the diseased parotid tissue. The extent of surgery may also depend on pathological examination of tissues removed during the surgery.

The nerve that controls motion to the face (the facial nerve) runs through the parotid gland. This nerve is important in closing the eyes, wrinkling the nose, and moving the lips. Most often the parotid gland can be removed without permanent damage to the nerve, however, the size and position of the diseased tissue may require that the nerve, or small branches of the nerve, be cut to assure complete removal. Even if the nerve is not permanently injured, there may be decreased motion of the facial muscles as the nerve recovers from the surgical procedure. If facial motion does not fully return your physician will discuss with you ways to rehabilitate facial movement.

Other possible short term complications include bleeding and infection. Although rare in parotid surgery, some patients may develop a thick scar or keloid. Many patients experience numbing of the earlobe and outer edge of the ear after parotid surgery. This generally resolves slowly over time. In a small proportion of patients the face on the side of the parotidectomy sweats at mealtimes, ( "gustatory sweating"). Most often this goes essentially unnoticed, however, if it should become bothersome medication and sometimes surgery are available.

Depending on the final diagnosis after the tissue is reviewed by a pathologist, additional diagnostic tests and follow-up examinations may be needed. Most often masses of the parotid are benign, and complete removal is the only treatment needed.

***Important Disclaimer Notice (Updated 8/7/14)***

Clinical indicators for otolaryngology serve as a checklist for practitioners and a quality care review tool for clinical departments. The American Academy of Otolaryngology—Head and Neck Surgery, Inc. and Foundation (AAO-HNS/F) Clinical Indicators are intended as ***suggestions, not rules***, and should be modified by users when deemed medically necessary. In no sense do they represent a standard of care. The applicability of an indicator for a procedure must be determined by the responsible physician in light of all the circumstances presented by the individual patient. Adherence to these clinical indicators will not ensure successful treatment in every situation. The AAO-HNS/F emphasizes that these clinical indicators should not be deemed inclusive of all proper treatment decisions or methods of care, nor exclusive of other treatment decisions or methods of care reasonably directed to obtaining the same results. The



**AMERICAN ACADEMY OF  
OTOLARYNGOLOGY–  
HEAD AND NECK SURGERY**

AAO-HNS/F is not responsible for treatment decisions or care provided by individual physicians. Clinical indicators are not intended to and should not be treated as legal, medical, or business advice.

CPT five-digit codes, nomenclature and other data are copyright 2009 American Medical Association. All Rights Reserved. No fee schedules, basic units, relative values or related listings are included in CPT. The AMA assumes no liability for the data contained herein.

© 2010 American Academy of Otolaryngology-Head and Neck Surgery. 1650 Diagonal Road, Alexandria, VA 22314.