

AAO21: Otitis Media with Effusion: Hearing Test for Chronic OME > 3 months

High Priority: No; N/A

Percentage of patients age 6 months to 12 years of age at the time of the visit with a diagnosis of otitis media with effusion including chronic serous, mucoid, or nonsuppurative otitis media with effusion of ≥3 months duration who had audiometry performed, ordered or who were referred for an audiometric evaluation.

Denominator:

All patients age 6 months through 12 years of age with a diagnosis of otitis media with effusion including chronic serous, mucoid, or nonsuppurative otitis media with effusion of ≥3 months duration.

Denominator Exclusions:

None

Denominator Exceptions:

Patient who had a hearing test within 6 months of the original diagnosis of otitis media with effusion including chronic serous, mucoid, or nonsuppurative otitis media with effusion.

Numerator:

Patients who had a hearing test performed, ordered or who were referred for an audiometric evaluation.

Measure Classifications:

Measure Type: Process

National Quality Strategy (NQS) domain: Effective Clinical Care

Meaningful Measure Area: Preventive Care

Calculation:

Inverse measure: No Continuous measure: No Proportional measure: Yes

Ratio measure: No

Risk Adjusted measure: No

 $@\ 2020\ American\ Academy\ of\ Otolaryngology-Head\ and\ Neck\ Surgery\ Foundation.\ All\ Rights\ Reserved.$

Limited proprietary coding is contained in the Measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets. The Measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, e.g. use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial uses of the Measures require a license agreement between the user and the American Academy of Otolaryngology—Head and Neck Surgery Foundation (AAO-HNSF).

Disclaimer: The Measures are not clinical guidelines, do not establish a standard of medical care, and have not been tested for all potential applications.

THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND. Neither the AAO-HNSF nor its members shall be responsible for any use of the Measures. The AAO-HNSF and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.

Empowering otolaryngologist-head and neck surgeons to deliver the best patient care