

AAO24: Allergic Rhinitis: Avoidance of Leukotriene Inhibitors

High Priority: Yes; Appropriate Use

Percentage of patients age 2 years and older patients with allergic rhinitis who do not receive leukotriene inhibitors.

Denominator:

All patients age 2 years and older with allergic rhinitis with a diagnosis of allergic rhinitis who do not have an active order (prescription or reported over the counter use) of intranasal corticosteroids or second generation antihistamines.

Denominator Exclusions:

Exclude patients with concomitant diagnosis of asthma or sleep apnea.

Denominator Exceptions:

Patients with allergy to intranasal corticosteroids, allergy to second generation antihistamines, or prostate issues (due to concern for intolerance of primary therapy options).

Numerator:

Patients who do not receive leukotriene inhibitors.

Measure Classifications:

Measure Type: Process

National Quality Strategy (NQS) domain: Effective Clinical Care

Meaningful Measure Area: Medication Management

Calculation: 1st Performance Rate Inverse measure: No Continuous measure: No Proportional measure: Yes

Ratio measure: No

Risk Adjusted measure: No

© 2020 American Academy of Otolaryngology–Head and Neck Surgery Foundation. All Rights Reserved.

Limited proprietary coding is contained in the Measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets. The Measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, e.g. use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial uses of the Measures require a license agreement between the user and the American Academy of Otolaryngology—Head and Neck Surgery Foundation (AAO-HNSF).

Disclaimer: The Measures are not clinical guidelines, do not establish a standard of medical care, and have not been tested for all potential applications.

THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND. Neither the AAO-HNSF nor its members shall be responsible for any use of the Measures. The AAO-HNSF and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.

Empowering otolaryngologist-head and neck surgeons to deliver the best patient care

AAO24

Allergic Rhinitis: Avoidance of Leukotriene Inhibitors


MEASURE DESCRIPTION

Percentage of patients with allergic rhinitis who do not receive leukotriene inhibitors.



DENOMINATOR

INITIAL PATIENT POPULATION (IPP)



NUMERATOR

