

AAO17: Age-related Hearing Loss: Advanced Diagnostic Imaging of Bilateral Presbycusis or Symmetric SNHL

High Priority: Yes; Appropriate Use

Percentage of patients age 60 years and older with a diagnosis of bilateral presbycusis or symmetric sensorineural hearing loss who were NOT ordered magnetic resonance imaging (MRI) or a computed tomography scan (CT scan) of the brain, temporal bone or internal auditory canal for the primary indication of hearing loss.

Denominator:

Patients age 60 years and older diagnosed with bilateral presbycusis or symmetric sensorineural hearing loss.

Denominator Exclusions:

None

Denominator Exceptions:

Medical reason that may require an MRI or CT scan of the brain, temporal bone or internal auditory canal, including:

- Evaluation for cochlear implantation or surgical management of hearing loss
- Unilateral or pulsatile tinnitus
- Vertigo, disequilibrium, dizziness
- Asymmetrical hearing loss
- Acquired hearing loss following meningitis, measles, mumps
- Chronic otitis media, otosclerosis
- Head injury or trauma

Numerator:

Patients who were NOT ordered magnetic resonance imaging (MRI) or a computed tomography scan (CT scan) of the brain, temporal bone or internal auditory canal for the primary indication of age-related hearing loss. This restriction is relevant at any time interval after the diagnosis of age-related hearing loss.

Measure Classifications:

Measure Type: Process

National Quality Strategy (NQS) domain: Efficiency and Cost Reduction

Meaningful Measure Area: Appropriate Use of Healthcare

Calculation: 1st Performance Rate

Inverse Measure: No

Continuous Measure: No

Proportional Measure: Yes

Ratio Measure: No

Risk Adjusted: No

Copyright:

© 2019 American Academy of Otolaryngology – Head and Neck Surgery Foundation. All Rights Reserved.

Limited proprietary coding is contained in the Measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets.

The Measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, e.g. use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain.

Commercial uses of the Measures require a license agreement between the user and the American Academy of Otolaryngology – Head and Neck Surgery Foundation (AAO-HNSF).

Empowering otolaryngologist–head and neck surgeons to deliver the best patient care

Disclaimer: The Measures are not clinical guidelines, do not establish a standard of medical care, and have not been tested for all potential applications.

THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND. Neither the AAO-HNSF nor its members shall be responsible for any use of the Measures. The AAO-HNSF and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.