

## AAO12: Tympanostomy Tubes: Topical Ear Drop Monotherapy for Acute Otorrhea

**High Priority Status:** Yes; Appropriate Use  
**NQF Number:** N/A

Percentage of patients age 6 months to 12 years of age at the time of the visit with a current diagnosis of an uncomplicated acute tympanostomy tube otorrhea (TTO) who were prescribed or recommended to use topical antibiotic eardrops and NOT prescribed systemic (IV or PO) antibiotics for acute TTO.

### Denominator:

All patients age 6 months through 12 years of age at the time of the visit with a current diagnosis of an uncomplicated\* acute\*\* TTO

\* Uncomplicated refers to TTO that is not accompanied by high fever (38.5°C, 101.3°F), concurrent illness requiring systemic antibiotics (e.g., streptococcal pharyngitis, bacterial sinusitis), or cellulitis extending beyond the external ear canal to involve the pinna or adjacent skin.

\*\* Acute refers to less than four weeks' duration.

### Denominator Exclusions:

None

### Denominator Exceptions:

Medical reason(s) for not prescribing/recommending topical antibiotic monotherapy or for prescribing oral antibiotics include:

- Patients with high fever;
- Patients with cellulitis of the ear or adjacent skin;
- Patients who are immunocompromised;
- Patients with oral antibiotic given for other concurrent condition including but not limited to streptococcal pharyngitis, bacterial sinusitis, pneumonia, pharyngitis, osteitis abscess, or other concurrent acute infection beyond the confines of the ear canal;
- Patients who failed topical antibiotic therapy;
- Patients with a cochlear implant;
- Patients with concurrent diabetes;
- Patients who were prescribed oral antibiotics by another clinician; concurrent diagnosis of acute suppurative otitis media with effusion with intact eardrum; acute TTO persists, or worsens, despite topical antibiotic therapy.

Patient reason(s) for not prescribing/recommending topical antibiotics or prescribing oral antibiotics include:

- Administration of eardrops is not possible because of local discomfort;
- Lack of tolerance.

### Numerator:

Patients who were prescribed or recommended to use topical antibiotic eardrops and NOT prescribed systemic (IV or PO) antibiotics for acute TTO.

### Numerator Exclusions:

None

### Measure Classifications:

Measure Type: Process

National Quality Strategy (NQS) domain: Effective Clinical Care

Meaningful Measures Area: Appropriate Use of Healthcare

Care Setting(s): Ambulatory Care: Clinician Office/Clinic

Includes Telehealth: Yes

Number of Performance Rates: 1

Inverse measure: No

Continuous measure: No

Proportional measure: Yes

Ratio measure: No

Risk-adjusted measure: No

© 2022 American Academy of Otolaryngology–Head and Neck Surgery Foundation. All rights reserved.


Limited proprietary coding is contained in the measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets. The measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, e.g. use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the measures for commercial gain, or incorporation of the measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial uses of the measures require a license agreement between the user and the American Academy of Otolaryngology – Head and Neck Surgery Foundation.

The measure is not a clinical guideline, does not establish a standard of medical care, and has not been tested for all potential applications. The measure and specifications are provided “as is” without warranty of any kind. Neither the American Academy of Otolaryngology – Head and Neck Surgery Foundation (AAO-HNSF), nor its members shall be responsible for any use of the measure. The AAO-HNSF and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.

### INITIAL PATIENT POPULATION (IPP)

#### MEASURE DESCRIPTION

Percentage of patients age 6 months to 12 years of age at the time of the visit with a current diagnosis of an uncomplicated acute tympanostomy tube otorrhea (TTO) who were prescribed or recommended to use topical antibiotic eardrops and NOT prescribed systemic (IV or PO) antibiotics for acute TTO.



### DENOMINATOR


#### \*Denominator Exceptions:

##### MEDICAL REASONS:

- Fever > 101.3° F
- Cellulitis of the ear
- Immunocompromised
- Oral antibiotics for concurrent condition
- Failed topical antibiotic therapy
- Cochlear implant
- Diabetic
- Prescribed antibiotics by another physician
- Concurrent diagnosis of acute suppurative otitis media
- Acute otorrhea persists or worsens

##### PATIENT REASONS:

- Local discomfort
- Intolerance for topical monotherapy



### NUMERATOR

