



AAO20: Tympanostomy Tubes: Hearing Test

High Priority Status: No; N/A
NQF Number: N/A

Percentage of patients age 6 months through 12 years with a diagnosis of otitis media with effusion (OME) who received tympanostomy tube insertion and received a hearing test within 6 months prior to tympanostomy tube insertion.

Denominator:

All patients age 6 months through 12 years with a diagnosis of OME who received tympanostomy tube insertion.

Denominator Exclusions:

None

Denominator Exceptions:

Patient or caregiver(s) refusal of hearing test.

Numerator:

Patients who had a hearing test performed within 6 months prior to tympanostomy tube insertion.

Numerator Exclusions:

None

Measure Classifications:

Measure Type: Process

National Quality Strategy (NQS) domain: Effective Clinical Care

Meaningful Measures Area: Appropriate Use of Healthcare

Care Setting(s): Ambulatory Care: Clinician Office/Clinic

Includes Telehealth: No

Number of Performance Rates: 1

Inverse measure: No

Continuous measure: No

Proportional measure: Yes

Ratio measure: No

Risk-adjusted measure: No

© 2022 American Academy of Otolaryngology–Head and Neck Surgery Foundation. All rights reserved.


Limited proprietary coding is contained in the measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets. The measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, e.g. use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the measures for commercial gain, or incorporation of the measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial uses of the measures require a license agreement between the user and the American Academy of Otolaryngology – Head and Neck Surgery Foundation.

The measure is not a clinical guideline, does not establish a standard of medical care, and has not been tested for all potential applications. The measure and specifications are provided “as is” without warranty of any kind. Neither the American Academy of Otolaryngology – Head and Neck Surgery Foundation (AAO-HNSF), nor its members shall be responsible for any use of the measure. The AAO-HNSF and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.

INITIAL PATIENT POPULATION (IPP)

MEASURE DESCRIPTION

Percentage of patients age 6 months through 12 years with a diagnosis of otitis media with effusion (OME) who received tympanostomy tube insertion and received a hearing test within 6 months prior to tympanostomy tube insertion




DENOMINATOR

***Denominator Exceptions:**

PATIENT REASON

- Refusal of hearing test



NUMERATOR

