



AAO36: Tympanostomy Tubes: Resolution of Otitis Media with Effusion (OME) in Adults and Children

High Priority Status: Yes; Outcome
NQF Number: N/A

Percentage of patients aged 6 months and older with a diagnosis of otitis media with effusion (OME) who are seen 2 to 8 weeks after tympanostomy tube surgery and OME is resolved.

Denominator:
Patients aged 6 months and older with a diagnosis of OME who are seen 2 to 8 weeks after tympanostomy tube surgery.

Denominator Exclusions:
None

Denominator Exceptions:
Patient refusal for follow-up examination.

Numerator:
Patients with resolution of OME.

Numerator Exclusions:
None

Measure Classifications:
Measure Type: Outcome
National Quality Strategy (NQS) domain: Effective Clinical Care
Meaningful Measures Area: Patient-Focused Episode of Care
Care Setting(s): Ambulatory Care: Clinician Office/Clinic
Includes Telehealth: Yes
Number of Performance Rates: 1
Inverse measure: No
Continuous measure: No
Proportional measure: Yes
Ratio measure: No
Risk-adjusted measure: No

© 2022 American Academy of Otolaryngology–Head and Neck Surgery Foundation. All rights reserved.

Limited proprietary coding is contained in the measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets. The measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, e.g. use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the measures for commercial gain, or incorporation of the measures into a product or service that is sold, licensed or distributed for commercial gain. Commercial uses of the measures require a license agreement between the user and the American Academy of Otolaryngology – Head and Neck Surgery Foundation.


The measure is not a clinical guideline, does not establish a standard of medical care, and has not been tested for all potential applications. The measure and specifications are provided “as is” without warranty of any kind. Neither the American Academy of Otolaryngology – Head and Neck Surgery Foundation (AAO-HNSF), nor its members shall be responsible for any use of the measure. The AAO-HNSF and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.

AAO36 Tympanostomy Tubes: Resolution of Otitis Media with Effusion in Adults and Children

INITIAL PATIENT POPULATION (IPP)

MEASURE DESCRIPTION

Percentage of patients aged 6 months and older with a diagnosis of otitis media with effusion who are seen 2 to 8 weeks after tympanostomy tube surgery and otitis media with effusion is resolved.




DENOMINATOR

*Denominator Exceptions:

PATIENT REASONS

- Refusal of follow-up examination



NUMERATOR

