

AAO36: Tympanostomy Tubes: Resolution of Otitis Media with Effusion in Adults and Children

High Priority: Yes; Outcome

Percentage of patients aged 6 months and older with a diagnosis of otitis media with effusion who are seen 2 to 8 weeks after tympanostomy tube surgery and otitis media with effusion is resolved.

Denominator:

Patients aged 6 months and older with a diagnosis of otitis media with effusion who are seen 2 to 8 weeks after tympanostomy tube surgery.

Stratified by age:

Criteria 1: Patients 6 months - 12 years of age

Criteria 2: Patients 13 years and older

Criteria 3: All patients 6 months and older

Denominator Exclusions:

None

Denominator Exceptions:

None

Numerator:

Patients with resolution of Otitis Media with Effusion.

Measure Classifications:

Measure Type: Outcome

National Quality Strategy (NQS) domain: Effective Clinical Care

Meaningful Measure Area: Patient-Focused Episode of Care

Calculation: 1st Performance Rate

Inverse measure: No

Continuous measure: No

Proportional measure: Yes

Ratio measure: No

Risk Adjusted measure: No

© 2020 American Academy of Neurology and American Academy of Otolaryngology–Head and Neck Surgery Foundation. All Rights Reserved.

Limited proprietary coding is contained in the Measure specifications for convenience. Users of the proprietary code sets should obtain all necessary licenses from the owners of these code sets.

The Measures, while copyrighted, can be reproduced and distributed, without modification, for noncommercial purposes, e.g. use by health care providers in connection with their practices. Commercial use is defined as the sale, license, or distribution of the Measures for commercial gain, or incorporation of the Measures into a product or service that is sold, licensed or distributed for commercial gain.

Commercial uses of the Measures require a license agreement between the user, the American Academy of Neurology (AAN) and the American Academy of Otolaryngology–Head and Neck Surgery Foundation (AAO-HNSF).

Disclaimer: The Measures are not clinical guidelines, do not establish a standard of medical care, and have not been tested for all potential applications.


THE MEASURES AND SPECIFICATIONS ARE PROVIDED "AS IS" WITHOUT WARRANTY OF ANY KIND. Neither the AAO-HNSF nor its members shall be responsible for any use of the Measures. The AAO-HNSF and its members disclaim all liability for use or accuracy of any Current Procedural Terminology (CPT®) or other coding contained in the specifications.

AAO36 Tympanostomy Tubes: Resolution of Otitis Media with Effusion in Adults and Children


INITIAL PATIENT POPULATION (IPP)

MEASURE DESCRIPTION

Percentage of patients aged 6 months and older with a diagnosis of otitis media with effusion who are seen 2 to 8 weeks after tympanostomy tube surgery and otitis media with effusion is resolved.



DENOMINATOR



NUMERATOR

